

UNCLASSIFIED

AD NUMBER
ADB256780
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies only; Proprietary Info.; Sep 99. Other requests shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott St., Fort Detrick, MD 21702-5012.
AUTHORITY
USAMRMC ltr, 23 Aug 2001

THIS PAGE IS UNCLASSIFIED

AD _____

Award Number: DAMD17-97-1-7278

TITLE: Ribozyme-Mediated Repair of Mutant P53 Transcripts in
Breast Cancer Cells

PRINCIPAL INVESTIGATOR: Bruce Sullenger, Ph.D.

CONTRACTING ORGANIZATION: Duke University
Durham, North Carolina 27710

REPORT DATE: September 1999

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Distribution authorized to U.S. Government agencies
only (proprietary information, September 1999). Other requests for this
document shall be referred to U.S. Army Medical Research and Materiel Command,
504 Scott Street, Fort Detrick, Maryland 21702-5012.

The views, opinions and/or findings contained in this report are
those of the author(s) and should not be construed as an official
Department of the Army position, policy or decision unless so
designated by other documentation.

20000809 111

NOTICE

USING GOVERNMENT DRAWINGS, SPECIFICATIONS, OR OTHER DATA INCLUDED IN THIS DOCUMENT FOR ANY PURPOSE OTHER THAN GOVERNMENT PROCUREMENT DOES NOT IN ANY WAY OBLIGATE THE U.S. GOVERNMENT. THE FACT THAT THE GOVERNMENT FORMULATED OR SUPPLIED THE DRAWINGS, SPECIFICATIONS, OR OTHER DATA DOES NOT LICENSE THE HOLDER OR ANY OTHER PERSON OR CORPORATION; OR CONVEY ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE, OR SELL ANY PATENTED INVENTION THAT MAY RELATE TO THEM.

LIMITED RIGHTS LEGEND

Award Number: DAMD17-97-1-7278

Organization: Duke University, Durham, NC 27710

Location of Limited Rights Data (Pages):

Those portions of the technical data contained in this report marked as limited rights data shall not, without the written permission of the above contractor, be (a) released or disclosed outside the government, (b) used by the Government for manufacture or, in the case of computer software documentation, for preparing the same or similar computer software, or (c) used by a party other than the Government, except that the Government may release or disclose technical data to persons outside the Government, or permit the use of technical data by such persons, if (i) such release, disclosure, or use is necessary for emergency repair or overhaul or (ii) is a release or disclosure of technical data (other than detailed manufacturing or process data) to, or use of such data by, a foreign government that is in the interest of the Government and is required for evaluational or informational purposes, provided in either case that such release, disclosure or use is made subject to a prohibition that the person to whom the data is released or disclosed may not further use, release or disclose such data, and the contractor or subcontractor or subcontractor asserting the restriction is notified of such release, disclosure or use. This legend, together with the indications of the portions of this data which are subject to such limitations, shall be included on any reproduction hereof which includes any part of the portions subject to such limitations.

THIS TECHNICAL REPORT HAS BEEN REVIEWED AND IS APPROVED FOR PUBLICATION.

Patricia M. Mobern

5/26/00

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE September 1999	3. REPORT TYPE AND DATES COVERED Annual (15 Aug 98 -14 Aug 99)	
4. TITLE AND SUBTITLE Ribozyme-Mediated Repair of Mutant P53 Transcripts in Breast Cancer Cells			5. FUNDING NUMBERS DAMD17-97-1-7278	
6. AUTHOR(S) Bruce Sullenger Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Duke University Durham, North Carolina 27710 e-mail: b.sullenger@cgct.duke.edu			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Distribution authorized to U.S. Government agencies only (proprietary information, September 1999). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.				12b. DISTRIBUTION CODE
13. ABSTRACT (Maximum 200 Words) This project explores the potential of a new and innovative approach to human gene therapy that may prove to be useful for the treatment or prevention of a range of genetic diseases including many types of cancer. We have previously demonstrated that a trans-splicing group I ribozyme can be employed to repair mutant transcripts in <i>E. coli</i> (Sullenger and Cech, <i>Nature</i> 1994) and mammalian cells (Jones et al., <i>Nature Medicine</i> 1996; Lan et al., <i>Science</i> 1998). Ribozyme-mediated repair of mutant mRNAs associated with a range of human diseases is now experimentally tractable, and we have begun to assess the therapeutic potential of this process for the repair of mutant transcripts implicated in the development and progression of breast cancer. Because mutation of the <i>p53</i> gene is the most common genetic change seen in a wide variety of malignancies including breast cancer, we have initially focused on repair of mutant <i>p53</i> transcripts. Toward this end, we have mapped the accessible sites for ribozyme binding on <i>p53</i> RNAs, and using this information we have generated ribozymes that can react with and repair mutant <i>p53</i> transcripts in breast cancer and other cell lines.				
14. SUBJECT TERMS RNA Repair, Trans-splicing Ribozyme, Gene Therapy, Suppressor Oncogene, p53			15. NUMBER OF PAGES 15	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Limited	

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)
Prescribed by ANSI Std. Z39-18
298-102

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

____ Where copyrighted material is quoted, permission has been obtained to use such material.

____ Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

____ Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

☒ In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

☒ For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

BA In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

BA In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

____ In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Bruce A. Sullivan

PI - Signature

9/6/99

Date

Research Grant

Sullenger, Bruce A.

4. Table of Contents

	Page Number
Front Cover	1
Report Documentation Page	2
Foreword	3
Table of Contents	4
Introduction	5
Body	5
Key Research Accomplishments	13
Reportable Outcomes	13
Conclusions	13
References	14

5. Introduction

The overall goal of this project is to develop a new and innovative approach to human gene therapy that is based upon ribozyme-mediated repair of mutant mRNAs (Sullenger and Cech, 1994; Sullenger and Cech, 1995; Sullenger, 1995). This approach may be especially useful for the treatment of cancer because it should restore the regulated expression of oncogenes. Previously, we described how a trans-splicing ribozyme can be employed to repair mutant mRNAs in *Escherichia coli* (Sullenger and Cech, 1994). Subsequently we demonstrated that trans-splicing can also proceed in mammalian cells (Jones et al., 1996). Finally, we recently demonstrated that ribozymes could be employed to repair sickle β -globin transcripts in erythrocyte precursors from patients with sickle cell disease (Lan et al., 1998).

Many mutant messenger RNAs from dominant and suppressor oncogenes have been identified that appear to be involved in tumorigenesis. Repair of any of these mRNAs may yield new insights into tumor development and treatment. However, we have initially focused upon the repair of mutant transcripts from the *p53* tumor suppressor gene. Much evidence indicates that loss of the *p53* protein is associated with neoplastic transformation. Mutant *p53* transcripts are frequently found in a range of primary human tumors and tumor cell lines including mammary carcinomas (Nigro et al., 1989; Bartek et al., 1990; Malkin 1994). Thus we have sought to develop ribozymes that can repair the mutant *p53* transcripts present in these tumor cells in hope of using such ribozymes to either revert the transformed phenotype of these cells or induce apoptosis. Toward this end the specific objectives of this research proposal have not changed and are:

- 1.) To determine which regions of mutant *p53* transcripts are accessible to trans-splicing ribozymes.
- 2.) To construct trans-splicing ribozymes that can repair mutant *p53* transcripts and test them in vitro.
- 3.) To evaluate ribozyme-mediated *p53* repair in tissue culture cells after transient transfection.
- 4.) To determine if the ribozyme can repair endogenous mutant *p53* transcripts in breast cancer cell lines harboring defective copies of the *p53* gene.
- 5.) To ascertain if repair of mutant *p53* transcripts renders breast cancer cells less tumorigenic.

6. Body

6.1. Overview

We have made significant progress toward fulfilling several of our specific aims since we received this grant on August 15th, 1997. One accomplishment is that we have determined which regions of the *p53* transcript are accessible to trans-splicing ribozymes. Such mapping has been


performed upon p53 transcripts generated by in vitro transcription and mutant p53 mRNAs present in total cellular RNA isolated from MDA-MB-231 mammary carcinoma cells. These results fulfill our first specific aim: to map mutant p53 transcripts. In addition, we constructed two trans-splicing group I ribozymes that can recognize two of the most accessible regions on the p53 mRNA and have tested the ability of this ribozyme to react with the intended nucleotide on the targeted p53 transcript. Our results demonstrate that such trans-splicing ribozymes can react with p53 transcripts with high fidelity in the test tube as well as in MDA-MB-231 breast cancer cells. More recently, we have constructed plasmids containing ribozyme expression cassettes and have been able to demonstrate that trans-splicing ribozymes can repair mutant p53 transcripts in transiently transfected Saos-2 human osteosarcoma cells. These results partially fulfill our specific aims #2, #3 and 4: to create trans-splicing ribozymes that can repair p53 transcripts and test their activity in test tubes and in cells.

6.2 Studies and Results on Objective #1: To determine which regions of mutant p53 transcripts are accessible to trans-splicing ribozymes (six months).

To ascertain which regions of the p53 transcript are accessible to ribozymes, we developed an RNA mapping strategy that is based on a trans-splicing ribozyme library and RNA tagging (Jones et al., 1996). To generate the mapping library, the guide sequence of the *Tetrahymena* group I trans-splicing ribozyme was randomized such that the 5' end of the RNAs in the library began with 5'-GNNNNN-3' where "G" represents guanine and "N" represents equal amounts of the 4 nucleotides. To map the p53 transcript in vitro, the mapping library was incubated under splicing conditions with either p53 transcripts generated by in vitro transcription using T7 RNA polymerase or with p53 transcripts present in total cellular RNA isolated from MDA-MB-231 breast cancer cells. To identify accessible uridine residues, the trans-splicing reaction products were reverse transcribed (RT) and amplified by the polymerase chain reaction (PCR) with primers specific for the ribozyme's 3' exon tag (Jones et al., 1996) and for the 5' end of the p53 target RNA. From this analysis, the uridines at positions 41, 65 and 384 of p53 RNA appeared to be particularly accessible in p53 transcripts regardless of whether the p53 transcripts were generated by in vitro transcription or isolated from MDA-MB-231 mammary carcinoma cells (Fig. 1). These mapping results, taken together with the fact that tumor cells often have mutations downstream of the nucleotide at position 360, encouraged us to focus on developing ribozymes that recognize the uridines present at positions 41 and 65 on the p53 mRNA. Thus, the internal guide sequence on the L-21 trans-splicing ribozyme was changed to 5'-GGAGGG-3' to generate a ribozyme, called Rib41, specific for site 41 and to 5'-GGGUCU-3' to generate a ribozyme, called Rib65, specific for site 65. In addition, inactive versions of these ribozymes, called Rib41d and Rib65d, which lack part of the catalytic core of the enzyme, were generated as controls (Sullenger and Cech, 1994).

Figure 1: Mapping Results of the Accessible Uridines on the p53 Transcript

reaction sites (nt)	number of clones	
	in vitro	cellular RNA from MDA-MB-231 cells
+24	49	-
+41	4	4
+65	2	2
+307	-	1
+332	-	1
+340	1	1
<hr/>		
+384	6	5
+389	-	1


hot spots of mutations in human cancers

6.3 Studies and Results on Objective #2: To construct trans-splicing ribozymes that can repair mutant *p53* transcripts and test them in vitro (six months).

Rib41 and Rib65 can trans-splice a 3' exon tag onto *p53* transcripts in vitro. The trans-splicing ribozymes, Rib41-3'tag and Rib65-3'tag, were incubated under splicing conditions with *p53* RNA generated by in vitro transcription or total RNA isolated from MBA-MB-231 cells. To determine if trans-splicing had occurred in any of the RNA samples, RT-PCR analyses were performed with one primer specific for the *p53* target RNA and the other primer specific for the 3' exon tag sequence. An amplified fragment of the expected size was generated from samples containing Rib41-3'tag and Rib65-3' Tag and either in vitro transcribed *p53* RNA or total RNA isolated from breast cancer cells (unpublished results). No such RT-PCR product was generated from samples that lacked a ribozyme or that contained the inactive versions of the ribozymes. Sequence analysis of the spliced products demonstrated that the ribozyme had reacted with the intended uridine. More recently, we have changed the 3'-exon attached to Rib41 so that it encodes the wild type sequence for the *p53* transcript. Using this ribozyme, we have now been able to demonstrate that this ribozyme can repair *p53* transcripts, that contain a point mutation at nucleotide 820, in the test tube (unpublished results).

6.4. Studies and Results on Objective #4: To determine if the ribozyme can repair endogenous mutant *p53* transcripts in breast cancer cell lines harboring defective copies of the *p53* gene (nine months).

To determine if Rib41-3'tag could react with mutant *p53* transcripts inside human mammary carcinoma cells, the ribozyme was introduced into MBA-MB-231 cells via liposome-mediated transfection. RT/PCR amplification generated a fragment of the expected size (87 base pairs) from the total RNA isolated from the MBA-MB-231 cells that had been transfected with the active ribozyme (Fig. 2). By contrast no such product was generated from RNA samples isolated from cells that were not transfected or were transfected with the inactive ribozyme, Rib41d-3'tag. When Rib41-3'tag was added to the RNA extraction buffer used to isolate total RNA from a sample of mock transfected MBA-MB-231 cells, no amplification product was generated (Fig. 2) suggesting that the observed trans-splicing products were generated inside the transfected cells and not during RNA analysis. From these results we conclude that Rib41 can react with *p53* transcripts inside human breast cancer cells. To confirm this result, we subcloned and sequenced a number of these PCR generated cDNAs shown in figure 2. The splice junctions on each of the 6 clones tested were identical and the sequence of a representative clone is shown in figure 3. This sequence analysis demonstrated that the ribozyme had reacted with the uridine at position 41 in the *p53* transcript inside of these cells with high fidelity because in all 6 clones sequenced the 3' exon tag had been attached onto the proper uridine.

Figure 2: Amplification of p53 derived trans-splicing products





Figure 3: Sequence analysis of trans-splicing reaction products



MDA-MB-231 cells

Figure 4: Ribozyme expression cassette design and evaluation






Figure 5: Trans-splicing mediated repair of p53 transcripts in Soas-2 cells.

A. Ribozyme 41



B. Ribozyme 65



6.5. Studies and Results on Objective #3: To evaluate ribozyme-mediated p53 repair in tissue culture cells after transient transfection. (nine months).

To determine if Rib41 and Rib65 could repair mutant p53 transcripts following transient transfection, we first cloned the ribozymes into expression cassettes containing the CMV immediate early promoter and the preproinsulin polyadenylation signal sequence (Fig. 4). A number of expression cassettes were constructed that contained a variety of guide sequence lengths as well as the presence or absence of the P10 region of the ribozyme. Each of these ribozyme constructs were individually cotransfected with an expression cassette containing a truncated version of the p53 gene into Saos-2 human osteosarcoma cells which do not naturally contain p53 derived transcripts. Total RNA was harvested from these cells and analyzed for the presence of the repaired p53 transcript by reverse transcription and polymerase chain reaction techniques. As shown in figure 5, repaired RNAs could only be detected in RNA samples from cells expressing the active ribozymes and the truncated p53 substrate transcript; whereas, no repaired p53 RNA was detected in inactive ribozyme transfected cells.

7. Key Research Accomplishments

- Mapped accessible regions for ribozyme binding on p53 transcript
- Demonstrated the trans-splicing ribozymes can react with mutant p53 transcripts in breast cancer cells.
- Demonstrated that RNA polymerase II expressed ribozymes can repair mutant p53 transcripts in Saos-2 cells.

8. Reportable Outcomes

None to date, but we intend on submitting a manuscript describing p53 RNA mapping and repair in the next few months.

9. Conclusions

In summary, we have made significant progress toward fulfilling several of our specific aims since we received the funding for this research proposal on August 15th, 1997. We have determined which regions of the p53 transcript are accessible to trans-splicing ribozymes using a novel RNA mapping strategy based upon a library of ribozymes and 3'-exon tagging. Such mapping has suggested that the nucleotides at positions 41 and 65 on the p53 transcript are particularly accessible to trans-splicing ribozymes. These results fulfill our first specific aim: to map the accessible regions on mutant p53 transcripts. In addition, we constructed two trans-splicing group I ribozymes that can recognize the uridines at positions 41 and 65 on the p53 mRNA and have tested the ability of these ribozymes to react with the intended nucleotide on the targeted p53 transcripts in the test tube and inside the human mammary carcinoma cell line MDA-MB-231. Our results demonstrate that these trans-splicing ribozymes can react with p53 transcripts with high fidelity in the test tube and that Rib41 can react with mutant p53 transcripts in breast cancer cells. More recently, we have generated a number

of ribozyme expression cassettes and have been able to demonstrate that the ribozyme can repair truncated p53 transcripts following expression in cotransfected Saos-2 osteosarcoma cells. These results partially fulfill specific aims #2, 3 and 4: to create trans-splicing ribozymes that can repair p53 transcripts and test their activity in test tubes and in cells. This next year, we plan to begin to assess what phenotypic effect(s) such p53 RNA repair has upon these breast cancer and other cells.

10. References

- Bartek, J., Iggo, R. and Lane, D. P. (1990). Genetic and immunochemical analysis of mutant p53 in human breast cancer cell lines. *Oncogene* 5, 893-899.
- Jones, J. T., Lee, S.-W. and Sullenger, B. A. (1996). Tagging ribozyme reaction sites to follow trans-splicing in mammalian cells. *Nature Medicine* 2, 643-648.
- Lan, N., Howery, R. P., Lee, S.-W., Smith, C. A. and Sullenger, B. A. (1998). Ribozyme-mediated repair of sickle b-globin mRNAs in erythrocyte precursors. *Science* 280, 1593-1596.
- Malkin, D. (1994). Germline p53 mutations and heritable cancer. *Ann. Rev. Genet.* 28, 443-465.
- Nigro, J. M. et al. (1989). Mutations in the p53 gene occur in diverse human tumour types. *Nature* 342, 705-708.
- Sullenger, B. A. and Cech, T. R. (1994). Ribozyme-mediated repair of defective mRNA by targeted trans-splicing. *Nature* 317, 619-622.
- Sullenger, B. A. and Cech, T. R. (1995). RNA repair: a new possibility for gene therapy. *Journal of NIH Research* 7, 46-47.
- Sullenger, B. A. (1995). Revising messages traveling along the cellular information superhighway. *Chemistry & Biology* 2, 249-253.



DEPARTMENT OF THE ARMY
US ARMY MEDICAL RESEARCH AND MATERIEL COMMAND
504 SCOTT STREET
FORT DETRICK, MARYLAND 21702-5012

REPLY TO
ATTENTION OF:

MCMR-RMI-S (70-1y)

23 Aug 01

MEMORANDUM FOR Administrator, Defense Technical Information
Center (DTIC-OCA), 8725 John J. Kingman Road, Fort Belvoir,
VA 22060-6218


SUBJECT: Request Change in Distribution Statement

1. The U.S. Army Medical Research and Materiel Command has reexamined the need for the limitation assigned to the technical reports listed at enclosure. Request the limited distribution statement for these reports be changed to "Approved for public release; distribution unlimited." These reports should be released to the National Technical Information Service.

2. Point of contact for this request is Ms. Judy Pawlus at DSN 343-7322 or by e-mail at judy.pawlus@det.amedd.army.mil.

FOR THE COMMANDER:

Encl


PHYLLIS M. RINEHART
Deputy Chief of Staff for
Information Management

Reports to be Downgraded to Unlimited Distribution

ADB241560	ADB253628	ADB249654	ADB263448
ADB251657	ADB257757	ADB264967	ADB245021
ADB263525	ADB264736	ADB247697	ADB264544
ADB222448	ADB255427	ADB263453	ADB254454
ADB234468	ADB264757	ADB243646	
ADB249596	ADB232924	ADB263428	
ADB263270	ADB232927	ADB240500	
ADB231841	ADB245382	ADB253090	
ADB239007	ADB258158	ADB265236	
ADB263737	ADB264506	ADB264610	
ADB239263	ADB243027	ADB251613	
ADB251995	ADB233334	ADB237451	
ADB233106	ADB242926	ADB249671	
ADB262619	ADB262637	ADB262475	
ADB233111	ADB251649	ADB264579	
ADB240497	ADB264549	ADB244768	
ADB257618	ADB248354	ADB258553	
ADB240496	ADB258768	ADB244278	
ADB233747	ADB247842	ADB257305	
ADB240160	ADB264611	ADB245442	
ADB258646	ADB244931	ADB256780	
ADB264626	ADB263444	ADB264797	